

ROKO® PROFESSIONAL WC TRIO

DESCRIPTION

Liquid for disinfection of food-contact surfaces in food businesses and disinfection of surfaces and sanitary facilities in healthcare institutions (outside the medical area); the product has bactericidal, fungicidal, mycobactericidal and virucidal properties Active substance: active chlorine released by sodium hypochlorite EC 231-668-3, CAS 7681-52-9, Content 4.5 g/100 g

COMPOUDS

chlorine-based bleaching compounds, anionic surfactants, non-ionic surfactants

TECHNICAL REQUIREMENTS

Appearance.....transparent or opalescent liquid

pH~12.5

Density at 20°C, g/cm³1.07

APPLICATION

Bactericidal and fungicidal effect: Disinfection: pour undiluted liquid onto the inside of toilet bowls, bathtubs and wash basins and rinse after 15 min. Follow the same procedure for surface disinfection. The product has a bactericidal effect after 5 minutes. Bactericidal and fungicidal action after 15 minutes. Disinfect food-contact surfaces with diluted product (50 ml of product dissolved in 5 L of water), duration of action 20-30 minutes. Then rinse abundantly with warm water for consumption. Mycobactericidal (mycobacteria) and virucidal (polio viruses, adeno type 5, mouse norovirus) effects: Disinfection - undiluted preparation: Precede the disinfection process by washing the surface. Under clean conditions the product is mycobactericidal and virucidal after 15 minutes. Under dirty conditions undiluted product is mycobactericidal and virucidal after 30 minutes. Disinfection - diluted formulation The

disinfection process should be preceded by washing of surfaces, materials. The product is mycobactericidal and virucidal at a concentration of 1.6 % by volume after 15 minutes. Disinfection in hospitals and other healthcare institutions is effective at a concentration of 33.3 % by volume recommended contact time of 15 minutes. Use in well-ventilated areas. Ventilation time after application 10-15 minutes.